

Bibliography

1. 52° Congresso Nazionale della Società Italiana di Reumatologia - RIMINI 25-28 NOVEMBRE 2015
L'uso dell'acido ialuronico nelle condropatie del ginocchio: Regenflex Starter contro Bioplus nelle patologie traumatiche I. Capparucci (1), C. Bartolucci, A. Federici, Valentini, V. Vita (1) Dipartimento di Scienze Biomolecolari Facoltà di Scienze Motorie Insegnamento di Medicina dello Sport Uniurb, Urbino (Italy).
2. VII CONGRESSO NAZIONALE SISMeS - Ricerca e Formazione applicate alle Scienze Motorie e sportive Padova, 2-4 Ottobre 2015
The use of hyaluronic acid in chondral disorders of the knee: Regenflex Starter versus Bioplus in traumatic diseases I. Capparucci, C. Bartolucci, A. Federici, M. Valentini, V. Vita Università degli Studi di Urbino.
3. 51° Congresso Nazionale SIR e 17° Congresso Nazionale CROI - 26-29 Novembre 2014 Rimini TRATTAMENTO INTRARTICOLARE CON ACIDO IALURONICO PER LA LIGAMENTIZZAZIONE IN TRAPIANTO OMOLOGO IN CHIRURGIA SOSTITUTIVA DELLO LCA - I. Capparucci, C. Bartolucci, I. Testa, V. Vita (Urbino, L'Aquila).
4. THE JOURNAL OF ORTHOPAEDICS TRAUMA SURGERY AND RELATED RESEARCH - The use of hyaluronic acid in chondral disorders of the knee: single-phase system vs. two-phase system. - I. Capparucci, C. Bartolucci, A. Federici, I. Testa, M. Valentini. (Urbino, L'Aquila)
5. Antalgic effect and clinical tolerability of hyaluronic acid in patients with degenerative diseases of knee cartilage: an outpatient treatment survey - Castellacci E.; Polieri T., *Drugs Exp Clin Res*, 2004; 30(2): 67-73.
6. The Safety and Efficacy of Intra-Articular Dual Molecular Weighted Hyaluronic Acid in the Treatment of Knee Osteoarthritis: the I.D.E.H.A. Study - Shen X.; Gatti R., *Orthop Rev (Pavia)*, 2013; 5(4): e33.
7. Efficacy and safety of a single intra-articular injection of non-animal stabilized hyaluronic acid (NASHA) in patients with osteoarthritis of the knee - Altman R.D.; Åkermark C.; Beaulieu A.D.; Schnitzer T., *Osteoarthritis Cartilage*, 2004; 12(8): 642- 9.
8. A randomized saline-controlled trial of NASHA hyaluronic acid for knee osteoarthritis - Arden N. K.; Åkermark C.; Andersson M.; Todman M.G.; Altman R.D., *Curr Med Res Opin*, 2014; 30(2): 279-86.
9. NASHA hyaluronic acid vs methylprednisolone for knee osteoarthritis: a prospective, multi-centre, randomized, non-inferiority trial - Leighton R.; Åkermark C.; Therrien R.; Richardson J.B.; Andersson M.; Todman M.G.; Arden N.K., *Osteoarthritis Cartilage*, 2014;22(1): 17-25.
10. Intra-articular hyaluronic acid injection versus oral non-steroidal anti inflammatory drug for the treatment of knee osteoarthritis: a multi-center, randomized, open-label, non-inferiority trial - Ishijima M.; Nakamura T.; Shimizu K.; Hayashi K.; Kikuchi H.; Soen S.; Omori G.; Yamashita T.; Uchio Y.; Chiba J.; Ideno Y.; Kubota M.; Kurosawa H.; Kaneko K., *Arthritis Research & Therapy*, 2014; 16(1): R18.
11. Chang SC, Hoang B, Thomas JT, et al: Cartilage-derived morphogenetic proteins: New members of the transforming growth factor-beta superfamily predominantly expressed in long bones during human embryonic development. *J Biol Chem* 1994; 269: 28227-28234.
12. Jackson DW, Felt JC, Song Y, Van Sickle DC, Simon TM: Restoration of large femoral trochlear sulcus articular cartilage lesions using a flowable polymer: An experimental study in sheep. *Trans Orthop Res Soc* 2000;25:670.
13. Altman RD. Status of hyaluronan supplementation therapy in osteoarthritis. *Curr Rheumatol Rep*. 2003 Feb;5(1):7-14.
Castellacci E. et al. Antalgic effect and clinical tolerability of hyaluronic acid in patients with degenerative diseases of knee cartilage: an outpatient treatment survey. *Drugs Exp Clin Res*. 2004;30(2):67-73
14. Coleman PJ, et al. Role of hyaluronan chain length in buffering interstitial flow across synovium in rabbits. *J Physiol*. 2000 Jul 15;526 Pt 2:425-34
15. Ghosh P, et al. Potential mechanism of action of intra-articular hyaluronan therapy in osteoarthritis: are the effects molecular weight dependent? *Semin Arthritis Rheum*. 2002 Aug;32(1):10-37.
16. Kelly MA, et al. Intra-articular hyaluronans in knee osteoarthritis: rationale and practical considerations. *Am J Orthop*. 2004 Feb;33(2 Suppl):15-22.
Michel BA, et al. In press.
17. Moreland LW. Intra-articular hyaluronan (hyaluronic acid) and hylans for the treatment of osteoarthritis: mechanisms of action. *Arthritis Res Ther*. 2003;5(2):54-67.
18. Moskowitz RW, et al. Understanding osteoarthritis of the knee-causes and effects. *Am J - Orthop*. 2004 Feb;33(2 Suppl):5-9
19. Moskowitz RW. Hyaluronic acid supplementation. *Curr Rheumatol Rep*. 2000 Dec;2(6):466-71.
20. Pleimann JH, et al. Viscosupplementation for the arthritic ankle. *Foot Ankle Clin*. 2002 Sep;7(3):489-94.
21. Lotz M, et al. Value of biomarkers in osteoarthritis: current status and perspectives. *Ann Rheum Dis*. 2013 Nov 1;72(11):1756-63.
22. Reijman M, Hazes JM, Bierma-Zeinstra SM, et al. A new marker for osteoarthritis: cross-sectional and longitudinal approach. *Arthritis Rheum* 2004;50:2471-8.
23. Larsson S, Englund M, Struglics A, et al. The association between changes in synovial fluid levels of ARGS-aggregan fragments, progression of radiographic osteoarthritis and self-reported outcomes: a cohort study. *Osteoarthritis Cartilage* 2012;20:388-95.
24. Ku JH, Lee CK, Joo BS, et al. Correlation of synovial fluid leptin concentrations with the severity of osteoarthritis. *Clin Rheumatol* 2009;28:1431-5.
25. Schaffler A, Ehling A, Neumann E, et al. Adipocytokines in synovial fluid. *JAMA* 2003;290:1709-10.

